

Quantitative Analysis of Redox-Sensitive Proteome with DIGE and ICAT

Cexiong Fu,[†] Jun Hu,[†] Tong Liu,[†] Tetsuro Ago,[‡] Junichi Sadoshima,[‡] and Hong Li^{*,†}

Center for Advanced Proteomics Research and Department of Biochemistry and Molecular Biology, UMDNJ-New Jersey Medical School Cancer Center, Newark, New Jersey 07103, and Cardiovascular Research Institute and Department of Cell Biology and Molecular Medicine, UMDNJ-New Jersey Medical School, Newark, New Jersey 07103

Received February 6, 2008

Oxidative modifications of protein thiols are important mechanisms for regulating protein functions. The present study aimed to compare the relative effectiveness of two thiol-specific quantitative proteomic techniques, difference gel electrophoresis (DIGE) and isotope coded affinity tag (ICAT), for the discovery of redox-sensitive proteins in heart tissues. We found that these two methods were largely complementary; each could be used to reveal a set of unique redox-sensitive proteins. Some of these proteins are low-abundant signaling proteins and membrane proteins. From DIGE analysis, we found that both NF- κ B-repressing protein and epoxide hydrolase were sensitive to H₂O₂ oxidation. In ICAT analysis, we found that specific cysteines within sarcoplasmic endoplasmic reticulum calcium ATPase 2 and voltage-dependent anion-selective channel protein 1 were sensitive to H₂O₂ oxidation. From these analyses, we conclude that both methods should be employed for proteome-wide studies, to maximize the possibility of identifying proteins containing redox-sensitive cysteinyl thiols in complex biological systems.

Keywords: oxidation • DIGE • ICAT • cysteine thiol • redox proteomics

Introduction

Oxidative modifications of protein thiols serve important roles in regulating cellular physiology. Under mild oxidative stress, reversible oxidation of selective protein residues may serve as redox sensors and signal transducers for conveying cellular anti-stress responses. When encountering severe oxidative insults, many proteins undergo irreversible oxidative modifications, causing protein degradation and cell death.¹ Among the many amino acids that are susceptible to oxidative stress, cysteine is particularly sensitive to oxidative insults, owing to its favorable nucleophilic property.² Given the unique chemical nature of protein thiols, they play essential roles in maintaining the integrity of protein structure and function. Redox-sensitive cysteines usually possess acidic pK_a's and are likely to deprotonate under physiological pH, rendering them susceptible to oxidant challenge. Redox-sensitive cysteines have been identified in a wide spectrum of proteins including transcription factors: OxyR,³ AP-1,⁴ NF- κ B,⁵ Ref-1,⁶ p53,⁷ and HIF-1 α ,⁸ signal transducers: cAMP-dependent protein kinase,⁹ mitogen-activated protein kinases,¹⁰ protein tyrosine phosphatase,¹¹ and apoptosis signal-regulating kinase 1;¹² and stress response proteins, such as peroxiredoxins,¹³ superoxide dis-

mutase,¹⁴ thioredoxin,¹⁵ and heat shock proteins.^{16,17} Oxidative modifications of these crucial cysteines have significant impacts on both protein structures and functions.

Reactive oxygen species (ROS) and reactive nitrogen species (RNS) are the major driving forces for oxidizing proteins in general and cysteine thiols in particular. ROS and RNS, including hydrogen peroxide (H₂O₂), superoxide ion (O₂^{•-}), nitric oxide (NO[•]), and hydroxyl radical (OH[•]), can be produced *in vivo* from a wide range of cellular processes, such as metabolism, proliferation, inflammation, and senescence. Certain reversible cysteine oxidative modifications, such as disulfide bond formation,¹⁸ sulfenation,^{19,20} nitrosylation,²¹ and glutathionylation,^{22,23} are suggested to serve cellular protective functions to preserve crucial cysteine residues from irreversible oxidative damage or to regulate redox signal transduction. On the other hand, refractory and irreversible oxidative modifications of cysteine residues such as sulfonic acid formation and 4-hydroxynonenal adducts formation often lead to the inhibition of protein function and/or protein degradation.¹

The labile, transient, and dynamic nature of reversible oxidative modifications poses enormous technical challenges for both accurate proteomic identification and sensitive quantification of cysteine thiols. Such difficulties also reside in the fact that many proteins of biological interest are present in relatively low abundance. A variety of gel-based proteomics methods have been developed to identify redox-sensitive proteins.^{24–27} Diagonal electrophoresis¹⁸ is designed to identify disulfide-bond interlaced proteins according to distinct gel

* To whom correspondence should be addressed. Hong Li, Department of Biochemistry and Molecular Biology, UMDNJ-NJMS, 185 S. Orange Ave. MSB E-609, Newark, NJ 07103. Tel: 973-972-8396. Fax: 973-972-5594. E-mail: liho2@umdnj.edu.

[†] Center for Advanced Proteomics Research.

[‡] Cardiovascular Research Institute.

migration profiles of proteins with either intra- or intermolecular disulfide bond linkages, under sequential nonreducing and reducing electrophoresis conditions. Protein glutathionylation and nitrosylation could be detected by 2DE methods by taking advantage of the availability of specific antibodies.²⁸ In addition, Kim et al.²⁷ devised a novel immunoblotting method based on the covalent attachment of a biotin-linked tag to reactive cysteines for immunoblotting detection. However, these immunoblotting methods suffer from moderate resolution, nonspecific detection, and low throughput. Recently, gel-based methods employing cysteine-specific fluorochromes^{24–26} or radioactive probes^{22,29} have been developed to identify redox-sensitive proteins within complex protein mixtures. To minimize gel-to-gel variations, it is essential to be able to carry out multiplexed experiments in a high-throughput fashion. Unlike conventional 2DE methods, the saturation-labeling DIGE method has the potential for robust multiplexed analysis of protein thiols. This approach employs a pair of fluorescent CyDyes to specifically label free thiols in multiple protein samples. Mixtures of differentially labeled proteins can then be resolved simultaneously in the same gel, and quantitative differences can be ascertained from the dual channel fluorescent images. Although DIGE was originally designed to measure the alterations of protein expression, Hurd and co-workers³⁰ have successfully tailored this technique to detect redox-sensitive proteins.

A gel-free mass spectrometry (MS)-based method, ICAT,³¹ is another thiol-specific proteomic technique with multiplexing capability. Each ICAT reagent consists of three essential groups: a thiol-reactive group, an isotope-coded light or heavy linker, and a biotin segment to facilitate peptide enrichment. In an ICAT experiment, protein samples are first labeled with either light or heavy ICAT reagents on cysteine thiols. The mixtures of labeled proteins are then digested by trypsin and separated through a multistep chromatographic separation procedure. Peptides are identified with tandem MS, and the relative quantifications of peptides are inferred from the integrated LC peak areas of the heavy and light versions of the ICAT-labeled peptides. Sethuraman et al.³² successfully applied the ICAT technique to discover 18 potential H₂O₂-sensitive membrane proteins in rabbit heart.

Despite reports of the successful applications of both DIGE and ICAT methods for the identification of redox-sensitive proteins, comparative studies of their strengths and limitations for analyzing redox proteomes have not been reported. Herein we used these two quantitative proteomic methods to screen for heart proteins that are sensitive to H₂O₂ oxidation. To our knowledge, this is the first direct comparison of DIGE and ICAT methods for the identification and quantification of redox-sensitive proteins in complex biological mixtures. We found that these two methods were largely complementary and each was able to reveal a set of unique low-abundant redox-sensitive proteins. From DIGE analysis, we were able to identify transcriptional regulators and signaling molecules that were H₂O₂-sensitive, whereas from ICAT analysis, we were able to discover several redox-sensitive membrane proteins. Based on this study, we conclude that both methods should be employed for redox proteomics studies in order to comprehensively identify redox-sensitive proteins as well as localize their reactive cysteine thiols within complex biological systems.

Materials and Methods

Chemicals and Reagents. HPLC grade acetonitrile (ACN) and water were purchased from J. T. Baker (Phillipsburg, NJ). Tris, α -cyano-4-hydroxycinnamic acid (CHCA), trifluoroacetic acid (TFA), iodoacetamide, dithiothreitol (DTT), and protease inhibitor cocktail were purchased from Sigma-Aldrich (St. Louis, MO). MS calibration standard peptides, Glu-fibrinopeptide, and human adrenocorticotrophic hormone 18–39 were also bought from Sigma-Aldrich. Cleavable ICAT reagent was obtained from Applied Biosystems (ABI, Forster city, CA). Porcine malate dehydrogenase (MDH) was purchased from Roche Applied Science (Indianapolis, IN).

Protein Extraction and Oxidation. Eight to ten week old Swiss Webster mice heart tissues (Pel Freeze Biologicals, Rogers, AR) were weighted (from four different animals, ~100 mg each) and lysed with 500 μ L of lysis buffer containing 8 M urea, 2% CHAPS, and 0.1% Triton X-100 in 30 mM tris at pH 8.0. A protease inhibitor cocktail was added prior to homogenization. The homogenates were centrifuged at 14 000g at 4 °C for 30 min. Protein concentrations in supernatants were measured by Bradford assay (Bio-Rad, Hercules, CA). Protein solution from each animal was divided into two aliquots of 500 μ g each; one was treated with 500 μ M H₂O₂ for 30 min in the dark, and the other was treated with an equal volume of H₂O. Protein samples were then desalted with a ReadyPrep 2D Cleanup Kit (Bio-Rad), and the protein pellets were reconstituted with either ICAT or DIGE labeling buffers in accordance to respective experiments.

DIGE Saturation Dye Labeling and 2DE. DIGE experiments were carried out in an octaplexed fashion, where Cy3 maleimide (Cy3m) was used to label the four control and four H₂O₂ treated samples. In parallel, Cy5 maleimide (Cy5m) was used to label eight equal aliquots of the internal standards. Each Cy3m-labeled sample was then mixed pairwise with one Cy5m-labeled internal standard and used for subsequent 2DE analysis. The DIGE labeling procedure was adapted from the manufacturer's protocol, except that the tris(2-carboxyethyl)phosphine (TCEP) reduction step was skipped to preserve oxidized cysteines. Briefly, protein samples were first resuspended in DIGE labeling buffer (8 M urea, 2% CHAPS, 0.1% Triton X-100, and 30 mM tris at pH 8.0). Five micrograms of proteins from either control or H₂O₂-treated samples was labeled with 4 nmol of Cy3m each. Aliquots of 5 μ g of the internal standards (prepared by pooling 20 μ g of proteins each from all eight samples) were labeled with 4 nmol of Cy5m separately. The labeling reactions were carried out at 37 °C for 1 h and then quenched by adding equal volumes of sample buffer (8 M urea, 2% CHAPS, 2% DTT, and 2% ampholytes 5–8). Each Cy5m-labeled internal standard was combined with a designated Cy3m-labeled sample (either control or H₂O₂ treated samples) for subsequent 2DE analysis. To achieve optimal DIGE labeling and prevent spontaneous thiol oxidations, all experiments were performed with minimal light exposure. In addition, each tube was flushed with nitrogen prior to labeling. The final volume of each combined reaction mixture was adjusted to 183 μ L with rehydration buffer (8 M urea, 2% CHAPS, 1% DTT, and 1% ampholytes 5–8) plus 2 μ L of 1% bromophenol blue. A pick gel sample was prepared by combining 25 μ g of proteins from all eight samples and paired with a Cy5m-labeled internal standard to facilitate spot matching. Samples were loaded onto 11 cm IPG strips (pH 5–8, Bio-Rad), rehydrated with the sample buffer for 2 h, and subsequently focused in a Protean IEF cell (Bio-Rad) for at least

80 000 V-hr, at 7000 V maximum and a current limit of 50 μ A/gel strip. The IEF-focused IPG strips were then sequentially equilibrated with 50 mM DTT and 50 mM iodoacetamide for 15 min each in equilibration buffer (6 M urea, 2% SDS, and 20% glycerol in 50 mM Tris at pH 8.3). Proteins were then separated in the second dimension with 12.5% Criterion Tris-HCl gels using a Criterion Doceca cell (Bio-Rad). After electrophoresis, gels were first rinsed with Milli-Q water and then fixed with 40% ethanol and 10% acetic acid. The gels were then scanned on a Typhoon 9400 imager (GE Healthcare, Piscataway, NJ) with 100 μ m resolution and appropriate photomultiplier tube voltages to ensure no spot saturation. The pick gel was stained overnight with Sypro Ruby (Bio-Rad) and scanned. DeCyder software (v5.0, GE Healthcare) was used to analyze the gel images. Gel images were processed with 1500 expected spots, and spots with areas smaller than 400 and slopes larger than 1.5, mostly dust artifacts, were excluded from further analysis. Changed protein spots with *p*-values ≤ 0.05 were matched to the pick gel and were excised for identification according to a protocol established in this laboratory.³³ Tryptic peptides were desalted with C₁₈ Ziptips (Millipore) and spotted onto a MALDI plate by mixing 1:1 with the matrix solution (6 mg/ml CHCA in 60% ACN and 0.1% TFA). MS and MS/MS spectra were acquired using a 4700 MALDI-TOF/TOF mass spectrometer (ABI). Protein identifications were performed by both peptide mass fingerprinting (with Profound, <http://prowl.rockefeller.edu/prowl-cgi/profound.exe>) and MS/MS spectra matching (using embedded Mascot search engine v1.9 on a GPS server v3.5, ABI). Database search parameters were as followed: mass tolerance of 50 ppm for MS and 0.3 Da for MS/MS, trypsin digestion with maximum one missed cleavage, NCBI mouse database, variable modifications including methionine oxidation, and carboamidomethylation of cysteines. Proteins identified with sequence coverage above 15% and at least two unique peptides identified with confidence interval (C. I.) values above 95% from the MS/MS search were considered significant.

ICAT Labeling and Multidimensional Chromatography. Protein samples were reconstituted with the labeling buffer supplied with the ICAT Reagent Kit (ABI) and processed according to the manufacturer's protocol, unless stated otherwise. Two ICAT labeling strategies were employed: (1) A forward strategy in which free cysteines were directly labeled in the control and treated samples with either light or heavy ICAT reagents respectively without the reduction of disulfide bonds. In brief, 100 μ g of the control sample was labeled with the light ICAT reagent, while the H₂O₂-treated sample was labeled with the heavy ICAT reagent. The TCEP reduction step was avoided to preserve the native thiol redox-states. The labeling reactions were carried out at 37 °C for 2 h, and excess ICAT reagents were quenched with additional 10 mM DTT. The newly reduced thiols were alkylated with 10 mM iodoacetamide. (2) A backward approach in which we first alkylated all free cysteine thiols with iodoacetamide and reduced the reversibly oxidized cysteines with TCEP. For easier conceptual comparison between the forward and backward labeling ICAT results, we used the light ICAT reagent to label the newly exposed cysteine thiols from H₂O₂-treated samples and heavy ICAT reagent for labeling the control samples. After the labeling reactions with either strategy, the light and heavy versions of the samples were combined and subjected to tryptic digestion at 50:1 (protein to enzyme) ratio overnight at 37 °C. Tryptic peptides were separated from the excess ICAT reagents and

detergents in the lysis buffer and were fractionated with strong cation exchange chromatography (SCX). SCX was carried out on a Biocad Sprint System (ABI) with a PolySulfoethyl A column (200 mm \times 4.6 mm, PolyLC Inc., Columbia, MD). The ICAT-labeled peptides were first acidified with mobile phase A (10 mM KH₂PO₄, 20% ACN at pH 2.7) to a final volume of 500 μ L and injected. The peptides were first separated at 1.0 mL/min with 0–25% mobile phase B (0.6 M KCl, 10 mM KH₂PO₄, and 20% ACN at pH 2.7) for 30 min, followed by 25–100% B over 20 min, and remained at 100% B for an additional 30 min. Peptide fractions were collected at 2 min intervals and dried. ICAT-labeled peptides were then enriched by avidin cartridges (ABI) and dried. Biotin moieties were cleaved off with TFA at 37 °C for 2 h. After the cleavage, the peptides were dried and reconstituted in 5% ACN and 0.1% TFA (mobile phase A for reversed phase LC, RPLC) and were purified with a LC Packings capillary HPLC system (Dionex, Sunnyvale, CA) on a PepMap C₁₈ column (5 μ m, 0.075 \times 150 mm) along with an inline trapping column (5 μ m, 0.3 \times 5 mm). RPLC gradient consisted of first 5–30% mobile phase B (MPB, 95% ACN, and 0.1% TFA) over 75 min, then 30–90% MPB for 15 min, and maintained at 90% MPB for an additional 10 min at a flow rate of 400 nL/min. The RPLC eluent was mixed at a 1:2 ratio with the MALDI matrix (6 mg/mL CHCA, 60% ACN, and 0.1% TFA) through a micro tee and spotted onto the MALDI plates with Probot (Dionex).³⁴

Redox Analysis of Purified Malate Dehydrogenase. One hundred micrograms of MDH was dissolved in 80 μ L of the lysis buffer (identical to the buffer used for heart protein extraction). Forty microliters each of MDH solution was treated with either H₂O or 500 μ M H₂O₂ for 30 min in the dark. Subsequently both DIGE and ICAT analyses were performed as described for the heart samples.

MS and Database Search. An ICAT-ratio-dependent acquisition strategy was used for the quantification and identification of peptides, where both MS and MS/MS spectra were acquired on a 4700 MALDI-TOF/TOF mass spectrometer (ABI). ICAT ion pairs (heavy and light versions with a mass difference of 9.03 \pm 0.03 Da) were quantified with the GPS Explorer software (v3.5, ABI). Relative ICAT ratios were computed with the integrated chromatographic areas of the ICAT ions. Only ICAT pairs with at least 20% intensity changes, plus with signal-to-noise ratios over 80 were submitted for MS/MS analysis and the subsequent MASCOT database search. Only peptides identified with C. I. values at or above 95% and mass errors within 20 ppm were considered as confident identifications. The frequency of false identification in ICAT analysis was evaluated with a target-decoy database search strategy proposed by Peng et al.³⁵ The decoy database was constructed with all database protein entry sequences reversed. The false positive rate (FPR) was calculated with the equation $FPR = (2 \times FP) / (FP + TP)$. False positive (FP) hit equals to the total number of peptide assignments in the decoy database search with C. I. $\geq 95\%$, and true positive (TP) hit is the number of total hits after the subtraction of FP hits. The FPR calculated for this study was 3.9%.

Results and Discussion

Establishment of a Model Redox System for the Comparison of ICAT and DIGE. We chose to use hydrogen peroxide treatment of extracted tissue proteins for this study was to create a robust model system with defined redox states for each protein, to compare the effectiveness of these two methods.

This model system is different from *in vivo* redox systems, in which many diverse and somewhat unpredictable oxidative modifications are likely to exist, rendering head-to-head comparisons of the ICAT and DIGE methods difficult. The H₂O₂ concentration selected was based on both our own observations and previous published reports.³⁶ We wanted to establish a model system that would demonstrate noticeable decrease of free cysteine thiols following oxidative treatment, yet not too severe to induce protein degradation. To correlate the relationship between the degree of protein thiol oxidation and H₂O₂ dosage, we conducted a Western blotting analysis to assess the extent of thiol alkylation changes with biotinylated iodoacetamide, following the oxidation of the heart proteins with serial-diluted H₂O₂. The same samples were also analyzed by 2DE to evaluate the degrees of protein degradation. To our surprise, from 50 to 400 μ M, no consistent and substantial protein thiol oxidation was observed (data not shown). Above 1 mM, we started to observe protein degradation (data not shown). It is known that H₂O₂ is not as potent as other agents that include hydroxyl radicals and peroxynitrite for protein oxidation, both *in vivo* and *in vitro*.³⁷ Additional reason why appreciable protein oxidation was observed only at 500 μ M of H₂O₂ might also be due to the fact that, a small tripeptide antioxidant, GSH could be present in the protein extracts, and it would quench the effects of H₂O₂ at lower oxidant concentrations. Given these observations, we chose 500 μ M of H₂O₂ for this methodology comparison study.

Identification of H₂O₂-Sensitive Heart Proteins by DIGE.

Saturation-labeling DIGE utilizes fluorescent CyDyes linked to a thiol-reactive maleimide group. Such that the Cy3m and Cy5m CyDyes can label free thiols in protein samples, facilitating simultaneous protein thiol measurement. Herein we used DIGE method to identify redox-sensitive heart proteins. Individual control and H₂O₂-treated samples were paired with a common internal standard for accurate gel-to-gel spot matching and quantification. Because we performed CyDye labeling reactions without protein reduction and alkylation, only free thiols should be labeled and their relative ratios measured. The gel images were analyzed with the DeCyder software and the quantification of free thiol levels in the samples was normalized to the pooled internal standards. Only significantly changed ($p \leq 0.05$) protein spots with at least 50% DIGE signal decrease were subjected to MS identification. Overall, 1000 proteins spots were detected and matched among the gels; 55 spots were excised for identification by MS. Forty-two spots containing 26 unique proteins were confidently identified (Supplemental Table 1, Supporting Information). An example of the comparison between either a control sample (green, Figure 1A) or a H₂O₂-treated sample (green, Figure 1B) and their respective internal standards (red) is shown in Figure 1. Among the proteins having significantly decreased thiol signals upon H₂O₂ treatment, an example of DIGE analysis of malate dehydrogenase is shown (Figure 1C–G). A treatment/control ratio of 0.26 (p -value < 0.05) was observed. We observed that several protein spots shifted to more acidic forms as a result of H₂O₂ treatment (Figure 1A and B), indicating the possible occurrence of acidic post-translational modifications such as the oxidation of cysteine thiols to sulfenic, sulfinic, or sulfonic acids. Recently, several studies have reported that key cysteines oxidized into these more acidic forms may serve as redox sensors for modulating protein functions.^{38,39} Given our experimental design, there may be two possibilities for the reduction of fluorescence intensities observed for H₂O₂-treated proteins:

H₂O₂-induced protein degradation or cysteine thiol oxidation. To distinguish these two possibilities, we also analyzed the samples by conventional 2DE and stained with Sypro Ruby. No significant changes at the protein expression levels were observed (data not shown). Therefore, it is likely that the selective reduction of thiol signals in H₂O₂-treated samples is primarily due to the oxidation of redox-sensitive cysteines.

We also attempted to locate the specific CyDye-labeled cysteines in these protein spots by searching for Cy3m or Cy5m as well as other possible oxidative modifications of protein thiols in both MS and MS/MS spectra. However, these efforts were not fruitful, probably due to (1) possible inefficient ionization of CyDye-labeled peptides, (2) poor recovery of large peptides from the in-gel digestion steps, and (3) the DTT reduction step during in-gel digestion may have reduced some oxidative modifications of cysteines. Although the exact sites and types of oxidative modifications could not be determined solely by the DIGE method, it did provide a panoramic view of proteins that are sensitive to H₂O₂ treatment. Many proteins found here have previously been reported as sensitive to redox modifications (Table 1). For example, many metabolic proteins, especially those involved in the glycolysis, pyruvate dehydrogenase complex, and the tricarboxylic acid cycle, were found to be susceptible to H₂O₂ oxidation. In addition, proteins essential for the mitochondrial electron transport chain also underwent oxidative modifications upon H₂O₂ treatment. Finally, we found a significant impact of H₂O₂ on antioxidant proteins, structural and signaling proteins as well. The widespread oxidation of proteins involved in such diverse biological processes suggested that oxidative stress may impact many cellular machineries within the cardiovascular system. Interestingly, a low-abundant nuclear protein, NF- κ B repressing factor, was found oxidized upon H₂O₂ treatment. The finding may suggest a novel mechanism for NF- κ B regulation. The known pathway for NF- κ B activation by oxidative stress is initiated via the oxidation-induced dissociation of NF- κ B and its inhibitor, I κ B in the cytoplasm. The activated NF- κ B will translocate to the nucleus and activate target gene transcription.^{5,40} Under normal physiological conditions, NF- κ B repressing factor is the guardian in the nucleus that prevents abnormal activation of NF- κ B by direct protein–protein interaction. The oxidation of the NF- κ B repressing factor may abolish its NF- κ B binding capability and pave the way for full activation of NF- κ B.

Identification of H₂O₂-Sensitive Heart Proteins by ICAT.

We adopted a forward ICAT labeling strategy previously reported by Sethuraman et al.⁴¹ to identify redox-sensitive protein thiols. Modifications (as described in the Materials and Methods) were made to enhance both protein digestion and downstream MS/MS peptide fragmentation efficiency. The identification of the H₂O₂-sensitive cysteine-containing peptides must satisfy the followed criteria: (i) contain at least one ICAT modified cysteine; (ii) at least 20% decrease in ICAT MS ion intensity following H₂O₂ treatment; (iii) C. I. $\geq 95\%$, and (iv) each peptide assigned to only one protein without redundancy.

Over 2000 ICAT pairs were observed and quantified by MS; the histogram of all ICAT ratios showed a normal distribution with a median of 0.98 (data not shown), suggesting no significant bias was introduced during sample preparation and labeling. To assess the degree of random analytical variations and to obtain a cutoff value of the ICAT ratio that represents significantly oxidized peptides, a preliminary ICAT experiment of bovine serum albumin (BSA) standards in three predefined




Figure 1. Identification of H_2O_2 -sensitive heart proteins with saturation-labeling DIGE. Mouse heart proteins were incubated with H_2O or $500 \mu M H_2O_2$ and labeled with Cy3m (green), each sample was mixed with an internal standard labeled with Cy5m (red). A mixture of a control sample and the internal standard was resolved in a 2D gel and the superimposed fluorescent image from the control and the standard proteome is shown in (A). The same procedure was applied to a H_2O_2 -treated sample and the gel image is shown in (B). More green spots were observed in the control sample (A) than the H_2O_2 treated sample (B), when both are compared to the Cy5m labeled internal standard. This observation indicates that fewer free thiols were available for Cy3m labeling in the H_2O_2 treated sample (B). The white boxes in panels (A) and (B) highlight a group of succinate dehydrogenase isoforms that were sensitive to H_2O_2 oxidation (less green spots in B and confirmed with DeCyder quantification analysis). It is also noticeable that the acidic end of this group of protein isoforms appeared more red in A and greener in B, suggesting the appearance of more acidic isoforms under H_2O_2 challenge. Protein spots were detected and matched among the gels with the assistance of preassigned landmark spots. Consistent protein spot detection and matching is demonstrated between the control (C) and the H_2O_2 -treated gels (D). The highlighted spots in panels (C) and (D) represent a protein (later identified as malate dehydrogenase) with significant decrease of fluorescent intensity following H_2O_2 treatment. This change is more strikingly illustrated in the 3D comparison of the spot volumes between a control (E) and a H_2O_2 -treated sample (F). Quantitative analysis was carried out after spot volume normalization with the internal standard. Statistical evaluation ($p < 0.005$) (G) confirmed the significant decrease of $\sim 70\%$ of cysteine thiol in malate dehydrogenase following H_2O_2 treatment.

Table 1. Redox-Sensitive Proteins Found in This Study Using the Forward Labeling Strategy^a

Protein	DIGE	ICAT	peptide sequence	Previously reported redox features	Ref
Aconitase 2	+	+	³⁷⁹ VGLIGSCTNSSYEDMGR ³⁹⁵	Sulfonation of Cys ³⁸⁵	18, 20, 48
Isocitrate dehydrogenase subunit alpha	+	+	³⁵¹ CSDFTTEEICR ³⁶⁰	Disulfide bond formation of unspecified cysteines.	18
Malate dehydrogenase, cytosolic	+	+	¹²⁶ VIVVGNPANTNCLTASK ¹⁴²	Nitrosylation of cys ¹³⁷	21
Malate dehydrogenase, mitochondrial	-	+	⁷⁹ GYLGPEQLPDC ⁹¹ ⁹² GCDVVVIPAGVPR ¹⁰⁴ ²⁰⁴ TIPLISQCTPK ²¹⁵ ²⁷⁰ EGVVECSFVQSK ²⁸¹ ²⁸² ETECTYFSTPLLLGK ²⁹⁶	Redox-sensitive cys ⁹³	20, 32, 49
Oxoglutarate dehydrogenase	+	-		Possible carbonylation, unspecified cysteine	50
Succinyl-CoA ligase subunit alpha	-	+	¹⁵⁴ LIGPNC*PGVINPGECK ¹⁶⁹	Glutathionylation of cys ⁸⁹	51
Succinate dehydrogenase complex, subunit A	+	+	⁷⁶ AAFGLSEAGFN ⁹² ⁶⁴⁸ TLNEADCATVPPAIR ⁶⁶²	Intermolecular disulfide bonds	18
Dihydroliponamide S-succinyltransferase	+	-		Possible carbonylation	18, 50
Pyruvate dehydrogenase complex E2	+	-		Possible carbonylation	52
Pyruvate dehydrogenase E1 component subunit beta	-	+	²⁵⁹ EGIECEVINLR ²⁶⁹	Sulfenation on unspecified cysteine	20
Aspartate aminotransferase, mitochondrial	-	+	⁹⁵ EYLPIGGLAEFC ¹⁰⁷ ¹⁸⁶ TCGDFDFSGALEDISK ²⁰⁰	Carbonylation	52
Acyl-CoA dehydrogenase, short chain	+	-		Intermolecular disulfide bonds	18, 30
Very-long-chain acyl-CoA dehydrogenase	+	-			
Long-chain specific acyl-CoA dehydrogenase	-	+	¹⁶⁶ CIGAIAMTEPGAGSDLQGV ¹⁸⁵ ³⁴⁶ AFVDSCLQLHETK ³⁵⁸ ³⁴⁴ VNQIGSVTESLQA ³⁵⁸		
Alpha enolase	-	+	¹⁷⁴ YASICQQNGIVIVEPEILPDGDHDLKR ²⁰¹	Glutathionylation	53
Fructose-bisphosphate aldolase C	-	+	¹⁴⁴ IVSNASC*TTNCLAPLAK ¹⁶⁰	Redox-sensitive Cys ¹⁷⁸	54
Glyceraldehyde-3-phosphate dehydrogenase	-	+	²³³ VPTPNVSVVDLTCR ²⁴⁶ ⁴⁴ NTGICTIGPASR ⁵⁶ ²⁰⁷ IYGGSVTGATCK ²¹⁹	Redox-sensitive cys ¹⁵⁰ and cys ¹⁵⁴ HNE modified cys ²⁴⁵	18, 21, 32, 55, 56
Pyruvate kinase 3	+	+		S-thiolation	49
Triosephosphate isomerase	-	+			
Creatine kinase, mitochondria	+	+	³¹¹ LGYILTQPSNLTGLR ³²⁶	Redox-sensitive cys ³¹⁷	18, 21, 32, 45

Table 1. Continued

Protein	DIGE	ICAT	peptide sequence	Previously reported redox features		Ref
				OXPHOS	Transport	
Lactate dehydrogenase 2, B chain	+	-	147 TTYAGNALCTVK ¹⁵⁸	Unspecified cysteine	20	
Electron transfer flavoprotein subunit alpha	+	+	397 NALVSHLDGTTTPVCEDIGR ⁴¹⁵			
Ubiquinol-cytochrome-c reductase complex core protein 1	+	+	256 VVEEDA VPGLTPCR ²⁶⁹ 448 YFYDQCPAVAGYGPIEQPLPDYNR ⁴⁷⁰	Unspecified cysteine	54	
COX VIb-1	-	+	29 NCWQNYLDFHR ³⁹ 48 GGDVSVCEWYRR ⁵⁹ 578 LQINAQNCVHCK ⁵⁸⁹	Carbonylation	52	
Electron transferring flavoprotein dehydrogenase	+	+		Unspecified cysteine	18, 52, 54	
NADH dehydrogenase	+	-		Unspecified cysteine	52	
Sodium/potassium-transporting ATPase subunit beta-1	-	+	205 YNPNVLPVQCTGK ²¹⁷	Unspecified cysteine	57	
Serotransferrin precursor	-	+	526 CAPNNKEEYNGYTGAFR ⁵⁴² 655 CFVKLPEGTTPEK ⁶⁶⁷	Unspecified cysteine	42	
SERCA 2	-	+	437 VGEATETALCLVEK ⁴⁵¹ 864 VSYQLSHFLOCK ⁸⁷⁶	Cys ⁸⁷⁵ redox state regulates calcium pump switch	32	
VDAC 1	-	+	134 EHINLGCDVDFDIAGPSIR ¹⁵² 238 YQVDPDACFSAK ²⁴⁹	Redox-sensitive cys ^{140, 245}	19	
Actin, alpha cardiac muscle 1	-	+	218 LCYVALDFENEMATAASSSLEK ²⁴⁰	Unspecified cysteine	19	
Actinin alpha 2	+	+	853 ELPPDQAQYCIK ⁸⁶⁴ 339 CQLEINFNTLQTK ³⁵¹	Unspecified cysteine	19	
Myosin-binding protein C, cardiac-type	+	+	416 TLTISQC SLADDAAYQC*VVGGEK ⁴³⁸	Unspecified cysteine	19	
Myosin light polypeptide 3	-	+	80 ITYGCQGDVLR ⁹⁰ 181 LMAGQEDSNGCINYEAFVK ¹⁹⁹	Unspecified cysteine	19, 20	
Troponin-1 alpha chain	-	+	190 CALEELKTVTNNLK ²⁰⁵	Unspecified cysteine	19	
60 kDa heat shock protein	-	+	431 AAVEEGIVLGGGCALLR ⁴⁴⁶	Unspecified cysteine	21, 58	
Peroxisome-5, mitochondrial precursor	-	+	83 GVLFGVPGAFTPGCSK ⁹⁸	Redox-sensitive Cys ⁴⁴²	43	
Superoxide dismutase 1	+	+	5 AVVLKGDGPVQGTIHFEQK ²⁴	Sulfenic acid or disulfide bond		
Rab GDP dissociation inhibitor beta	+	+	194 TDDYLDQPC*ETINR ²⁰⁸			
Transcription factor NRF	+	-				

^a TCA cycle, Tricarboxylic acid cycle; PDHC, Pyruvate dehydrogenase complex; OXPHOS, Oxidative phosphorylation.




Figure 2. Example of ICAT identification of H₂O₂-sensitive heart proteins. A MS/MS spectrum for a peptide (²⁰⁴TIPLISQCTPK²¹⁵) from MDH isolated from mouse heart is presented with a continuous series of y-ions for confident identification. The observation of a mass difference of 339 Da between y³ and y⁴ ions matched the mass of a heavy ICAT reagent-modified cysteine. The decrease of cysteine thiol was observed in the MS spectrum (see insert). The hallmark of a pair of the heavy and light ICAT-labeled peptides is a mass difference of 9.03 Da between the two adjacent monoisotopic peaks in the MS spectrum. The relative ICAT ratio for this peptide was 0.73, indicating ~27% loss of the redox-sensitive cysteine thiols upon H₂O₂ treatment.

heavy-to-light ICAT ratios was performed. We found that 99% of the sample populations were located within two standard deviations (~7%) from the expected values (Supplemental Figure 1, Supporting Information). Therefore, at a stringent cutoff value of 20% ICAT ratio alteration, it was unlikely that the changes were due to random analytical variations. About 300 ICAT ion pairs with at least 20% changes were subjected to downstream MS/MS identification. In total, 71 ICAT-labeled peptides from 50 H₂O₂-sensitive proteins were identified, out of which 15 were identified with at least 2 peptides (Supplemental Table 2, Supporting Information). A large number of these proteins are metabolic enzymes, components of the mitochondrial electron transport chain, antioxidants and stress response proteins, as well as structural and signaling proteins. A MS/MS spectrum of a peptide (²⁰⁴TIPLISQCTPK²¹⁵) from MDH is presented with a rather complete series of y-ions for confident identification (Figure 2). The observation of a mass difference of 339 Da between y³ and y⁴ ions matched the mass of a heavy ICAT reagent modified cysteine. Quantification of thiol levels was carried out at MS level (see insert in Figure 2). The hallmark of a pair of the heavy and light ICAT-labeled peptides is a mass difference of 9.03 Da between the two adjacent monoisotopic peaks. The relative ICAT ratio for this peptide was 0.73, indicating ~27% loss of the redox-sensitive

cysteine thiols upon H₂O₂ treatment. This is in contrast with ~74% loss of redox sensitive cysteine thiols found in DIGE analysis (Figure 1C–G), perhaps due to the fact that many other cysteines within MDH may also be oxidized^{21,32} (see later discussion). Many proteins found in our ICAT analysis have previously been identified as redox-sensitive, either with specified or unspecified redox-reactive cysteines (Table 1). Sarco-plasmic/endoplasmic reticulum calcium ATPase 2 (SERCA2) is an important transmembrane calcium pump. The SERCA2 pump machinery involves the formation of a redox-regulated disulfide bond that gates the influx of calcium flow as well as oscillation frequency. A site-directed mutagenesis study⁴² has identified this pair of crucial cysteines (cys⁸⁷⁵ and cys⁸⁸⁷) and the formation of the disulfide bond has been shown to be modulated by endoplasmic reticulum (ER) protein disulfide isomerase. We found that the redox-sensitive cys⁸⁷⁵ in peptide ⁸⁶⁴VSFYQLSHFLQCK⁸⁷⁶ (C: ICAT labeled) with a 50% decline of free thiol levels upon H₂O₂ treatment. The disulfide bond counterpart Cys⁸⁸⁷ resides in a tryptic peptide spanning 47 amino acid residues, which might be too large for efficient MS/MS fragmentation. Peroxiredoxins (PRX) are important antioxidant proteins that serve to detoxify peroxides. Unlike other 2-cys motif-containing PRXs, PRX5 contains only one conserved cys⁹⁶ at its redox-sensitive catalytic center,⁴³ which may form

an intramolecular disulfide bond with cys²⁰⁰. We identified the redox-sensitive cys⁹⁶ in the ICAT-labeled peptide ⁸³GVLFGVP-GAFTPGCSK⁹⁸ with a 40% decline of free thiol levels upon H₂O₂ treatment. To the best of our knowledge, this study represents the first unambiguous identification of this redox-sensitive cysteine in PRX5 by a proteomic method. Many proteins identified here, contain reactive cysteines concurring with published reports (Table 1). They included aconitase 2 (cys³⁸⁵), malate dehydrogenase (cys¹³⁷), fructose-bisphosphate aldolase C (cys¹⁷⁸), creatine kinase (cys³¹⁷), and HSP 60 (cys⁴⁴²). Redox modifications of these crucial cysteines lead to the modifications of protein activities. Some redox-sensitive cysteine thiols are possibly reported here for the first time, including cytochrome C oxidase VIb-1, sodium/potassium transporting ATPase β -1, tropomyosin-1, and actinin α -2.

Comparing our ICAT study results with the observations made by Sethuraman et al.,³² both studies utilized a similar “forward” labeling strategy and have demonstrated the effectiveness of the ICAT approach for redox studies. Sethuraman et al. focused on membrane proteins. Interestingly, several identical peptides were seen in both studies, including peptides from creatine kinase, malate dehydrogenase, GAPDH, VDACL1, and SERCA2, suggesting that the ICAT approach is robust for the detection of redox-sensitive proteins. Furthermore, the degree of oxidation of these cysteines is comparable despite the fact that we used only one-fourth of the H₂O₂ concentration used in Sethuraman’s work. For example, creatine kinase was identified with 2 peptides, ¹⁷³GLSLPPACSR¹⁸² (9.8% oxidized) and ³¹¹LGYILTCPSNLGTGLR³²⁶ (42.3% oxidized) in their study and 8.0% and 37.0% in our report, respectively. Both studies demonstrated that cys³¹⁷ was more sensitive to H₂O₂ oxidation than cys¹⁸⁰. It is notable that the ¹⁷³GLSLPPACSR¹⁸² peptide is not included in Supplemental Table 2 (Supporting Information) in this study, due to the requirement of at least a 20% decrease in ICAT signals.

In addition to identifying diminished free thiol levels upon H₂O₂ treatment, we also attempted to identify other reversibly oxidized cysteines by including a DTT reduction step after ICAT labeling, followed by alkylation with iodoacetamide. Although in the present ICAT workflow, the opportunities to locate such cysteines were limited by the requirement of the coexistence of at least one ICAT-labeled cysteine in the same tryptic peptide; otherwise those carboamidomethyl modified cysteines would have been washed off during the avidin affinity enrichment step. We did identify 6 proteins containing carboamidomethyl modifications, including GAPDH, in which cys¹⁵⁰ has previously been shown to be essential for its function and the reversible oxidation of cys¹⁵⁰ could down-regulate GAPDH activity.⁴⁴ In our study, we observed a carboamidomethyl modified cys¹⁵⁰ together with an adjacent ICAT-labeled cys¹⁵⁴ in peptide IVSNASC¹⁵⁰TTNC¹⁵⁴LAPLAK. It is possible that GAPDH underwent reversible cysteine modification (e.g., sulfenation) to conserve the crucial cys¹⁵⁰ under H₂O₂ challenge. Despite the exact identity of the reversible modification could not be revealed by our current method, a modified method with the inclusion of different reducing reagents, such as arsenite (for sulfenic acid)²⁰ or ascorbic acid (for nitrosylation)²¹ could be used to identify such modifications. In summary, we have shown that ICAT is a versatile and robust method to

identify redox-sensitive proteins with precise specification of sites of oxidation.

Comparison of Forward and Backward Labeling Strategies.

Recently, Hurd and co-workers³⁰ reported a “backward” labeling strategy to identify redox-sensitive proteins with the DIGE method. With this strategy, free thiols were initially blocked by alkylating reagents, and those reversibly oxidized thiols were then reduced by DTT and labeled with different CyDyes. In contrast to this “backward” labeling strategy, we used a “forward” DIGE labeling strategy in this study, where oxidized thiols were maintained in their native states, and free thiols were labeled with different saturation CyDyes for the quantification of redox-sensitive protein thiols. Both strategies can identify redox-sensitive proteins, but with their unique pros and cons. For example, the forward strategy could be affected by spontaneous cysteine oxidation during sample extraction and analysis. Such oxidation events could be minimized with low temperature, short processing time, and inert gas protection. The backward strategy could be impeded by tedious reagent cleanup steps prior to the DIGE labeling, which is not desirable for routine quantitative study.

Interestingly, despite using different DIGE labeling approaches to unravel redox-sensitive proteins in mouse heart following H₂O₂ treatments, a comparable number of redox-sensitive proteins were identified in both our and Hurd et al.’s work. Several proteins were identified by both strategies; for example, oxoglutarate dehydrogenase, succinate dehydrogenase, isocitrate dehydrogenase, electron transfer flavoprotein, acyl-CoA dehydrogenase, and superoxide dismutase (see Supplemental Table 1, Supporting Information). On the other hand, each method also identified a unique set of proteins. This observation may in part be due to the different isoelectric focusing pI ranges used (pI 5–8 in our study and pI 3–10 in Hurd’s study). Curiously, several proteins from the mitochondrial electron transport chain as well as cardiac actin and aconitase 2 were not seen as sensitive to H₂O₂ treatment by Hurd et al. However, we observed significant reduction of thiol levels in these proteins (either by DIGE or ICAT analysis), which are in agreement with published reports of having redox-sensitive cysteines within these proteins.^{20,32} The reason that these proteins were not detected by the “backward” DIGE labeling strategy is unclear. It is notable that several basic and/or hydrophobic proteins, for example, creatine kinase and voltage-dependent anion channel 1 (VDACL1), were observed by Hurd et al., but not by our forward DIGE strategy. We did identify these proteins as H₂O₂-sensitive proteins, however, with the “forward” labeling strategy employed for the ICAT analysis.

One possible advantage of the backward labeling strategy is the likelihood of revealing redox-sensitive cysteines buried inside the proteins that may not be accessible by the bulky DIGE or ICAT reagents using the forward strategy. To verify this possibility, we conducted a comparative ICAT study between the forward and backward labeling strategies. It was apparent that using the backward ICAT labeling strategy, we found 35% more peptides sensitive to H₂O₂ oxidation than using the forward labeling strategy (Supplemental Tables 2 and 3, Supporting Information). This result may be attributed to the reduction of the disulfide bonds, after which cysteines became more accessible for subsequent ICAT labeling reagents. Twenty-seven peptides (in 22 unique proteins) were found to be H₂O₂-sensitive by both strategies. The common proteins found with both approaches included peptides within aspartate




Figure 3. Venn diagram of H_2O_2 -sensitive proteins discovered by DIGE and ICAT methods. We identified 50 proteins as potential targets of H_2O_2 oxidation by the ICAT method and 26 with the DIGE method. Of these, 13 proteins were identified with both methods. Overall, the ICAT technique enabled us to identify 24 more redox-sensitive proteins than the DIGE method.

aminotransferase, creatine kinase, PRX5, and triosephosphate isomerase etc. (Supplemental Tables 2 and 3, Supporting Information). Similar to the comparison between forward and backward DIGE strategies, each ICAT labeling strategy also enabled the discovery of a distinct list of nonoverlapping peptides, implying the complementary nature of these two approaches.

Comparison of DIGE and ICAT Methods for Thiol Oxidation Studies. In this study, we have compared two thiol-specific quantitative proteomic techniques for redox-sensitive protein identification. DIGE is a gel-base method and quantifies overall protein thiol levels whereas ICAT utilizes multidimensional chromatographic methods to selectively enrich and quantify isotope-encoded cysteine-containing peptides. These two methods appear to be complementary for the purpose of identifying redox-sensitive proteins (Figure 3). There were 13 proteins commonly found by both methods (Table 1). They included actinin 2, creatine kinase, aconitase 2, malate dehydrogenase, ubiquinol-cytochrome-c reductase complex core protein 1, and superoxide dismutase. Many of these proteins are known to be susceptible to oxidative stress in cells.^{30,45–47} There were 37 redox-sensitive proteins identified solely by ICAT method. We found that some of these proteins are hydrophobic membrane proteins, which may be poorly resolved in 2DE. Examples of these proteins included SERCA2 and membrane glycoprotein gp42. Another scenario for proteins detected only with ICAT may attribute to the fact that protein *pI* values exceed the isoelectric focusing range of 2DE (*pI* 5–8 used in this study). PRX5 (*pI* 9.1), succinyl-CoA ligase (*pI* 9.46), cytochrome c oxidase VIb-1 (*pI* 8.96), CRP2 (*pI* 8.96), CRP3 (*pI* 8.90), and tropomyosin-1 alpha chain (*pI* 4.69) all fell into this category. Additional possibilities for ICAT-identified proteins that were missed in DIGE analysis might be due to the comigration of low and high abundance proteins to the same gel spots. It is conceivable that redox-sensitive proteins may not be identified in the background of abundant proteins. On the other hand, DIGE method also captured 13 proteins that were absent from ICAT analysis (Figure 3). They included oxoglutarate dehydrogenase, dihydrolipoamide S-acetyltransferase, epoxide hydrolase, NF- κ B-repressing factor, etc. (Supplemental Table 1, Supporting Information). We speculate that the lack of detection of these proteins with ICAT could result from incomplete enzymatic digestion, weak retentions on the RPLC trapping column, or poor ionization and/or fragmentation in MS. Furthermore, we compared the quantitative aspects of these two methods with regard to the redox-sensitive proteins found within the TCA cycle (Figure 4). Both methods found that

aconitase, isocitrate dehydrogenase, succinate dehydrogenase, and malate dehydrogenase were significantly oxidized upon H_2O_2 challenge. The ICAT method detected thiol reductions of 20–30% whereas the DIGE method observed more dramatic changes ranging from 70 to 90% of the thiol signals for the same proteins. On the other hand, succinyl-CoA ligase was observed solely in ICAT method with a 35% decrease, whereas oxoglutarate dehydrogenase oxidation appeared only in DIGE analysis with an over 95% decrease of free thiol signal. Overall, the trend of protein/peptide thiol signal decreases followed the same direction but to different extents. Almost in all cases, DIGE analysis reported more significant reduction of free thiol levels whereas ICAT only detected a moderate change. This is probably due to the fact that DIGE signals represented the quantification of a specific protein isoform that may differ from other isoforms of the same protein by having multiple oxidatively modified cysteines. In contrast, ICAT analysis detected the averaged changes of a specific cysteine residue among all protein isoforms.

To confirm the hypothesis that the quantitative discrepancies between DIGE and ICAT originated from the formation of a series of acidic protein isoforms following H_2O_2 oxidation, we have performed additional ICAT and DIGE analyses of a purified malate dehydrogenase (MDH) following identical H_2O_2 oxidation treatment used for heart proteins. From DIGE analysis, we found clear evidence of the appearance of additional acidic MDH isoforms (Figure 5, red gel spots 1–3) in H_2O_2 -treated MDH. The 7 discrete 2DE spots appeared to represent MDH isoforms with different extent of oxidatively acidified cysteines. The most acidic protein spots were likely to contain many but not all of the 8 MDH cysteines as sulfinic, sulfenic, or sulfonic acids, with the remaining cysteine thiols labeled with CyDyes. The DIGE ratios shown in Figure 5 were indicative of the relative amounts of the remaining cysteine thiol levels, likely barometers of the relative proportion of each MDH isoform in the samples under comparison, assuming overall MDH levels remained the same following H_2O_2 oxidation. The conclusions that can be drawn from DIGE analysis of MDH included that higher proportion of acidically oxidized MDH isoforms were formed following H_2O_2 oxidation. It should be mentioned that completely oxidized protein isoform would not be labeled by CyDyes but may be revealed with Sypro Ruby staining of the 2D gels. In ICAT experiments, we successfully identified 5 cysteine-containing MDH peptides, with the remaining peptide too large to be efficiently fragmented by MS/MS in our MS instrument. The quantification of these ICAT-labeled MDH peptides demonstrated varied susceptibility to H_2O_2 oxidation among the 5 cysteines (Table 2 and Supplemental Figure 2, Supporting Information). As a simplistic interpretation, Cys 69 was the most vulnerable to H_2O_2 oxidation, with 77% of its free thiol oxidized by H_2O_2 . Cys 65 was also relatively sensitive to H_2O_2 oxidation, with over 66% of free thiol oxidized. Cys 188, Cys 251, and Cys 261 were only moderately sensitive to H_2O_2 oxidation (Table 2). Both DIGE and ICAT redox quantification results should be interpreted with the understanding that related yet different biochemical information are obtained with each technique. The quantification results from DIGE method is gel spot-based (or protein isoform-specific) instead of simply protein-specific. From DIGE analysis, we found the increase of MDH oxidative isoforms. It was an effective method for discovering a small fraction of MDH isoforms that were particularly sensitive to redox regulation. On the other hand, with the ICAT method, the quantifica-




Figure 4. H_2O_2 -sensitive TCA cycle proteins identified in this study. Both ICAT and DIGE methods revealed that many proteins in the TCA cycle were prone to oxidation by H_2O_2 . The quantification of protein thiol level change with the ICAT method is the average of all its ICAT peptides. We detected a reduction of 20–30% of free cysteine thiols in aconitase 2, isocitrate dehydrogenase, succinate dehydrogenase, and malate dehydrogenase with the ICAT method, whereas a more dramatic reduction ranging from 70 to 90% for the same proteins was seen with the DIGE method. Succinyl-CoA ligase was observed solely by the ICAT method with a 35% free thiol level decrease, whereas oxoglutarate dehydrogenase oxidation was observed only with the DIGE method, with an over 95% decrease of free thiols. Student *t* test was carried out with four independent studies, *, $p < 0.05$; ***, $p < 0.005$.

tion is carried out at the peptide level, which is the average of all common peptides derived from various isoforms of the same protein. For the five peptides detected from MDH, the reported ICAT peptide ratios represented the relative amounts of total free thiols of the given peptides derived from the 7 isoforms observed in the DIGE gel. Therefore, potentially pronounced changes of a peptide redox state in one isoform may be “averaged out” by a large proportion of the same peptides present in the less oxidized isoforms, resulting in the overall marginal change reported by the ICAT ratios. Consequently,

only the redox trend can be compared between the two methods, indicating overall protein susceptibility to oxidative modifications. Because gel spots representing dramatically oxidized protein isoforms are more easily detected and quantified by DIGE software, it is therefore likely that the degree of oxidation reported from DIGE is more pronounced than ICAT, as reported in this study.

Adaptation of DIGE and ICAT Methods for Redox Proteomics Studies of Biological Systems. We chose denaturing buffer for this study to recover sufficient low-abundant proteins




Figure 5. DIGE analysis of MDH oxidized by H₂O₂. Cy3m (green)-labeled control sample and Cy5m (red)-labeled H₂O₂-treated porcine MDH. A series of MDH isoforms were observed with both untreated and H₂O₂-treated MDH. It was clear that the H₂O₂-treated MDH contained additional isoforms distributed toward the acidic end of the 2DE gel (red spots 1–3), suggesting the acidification of MDH, likely in the forms of cysteine oxidation to sulfenic, sulfenic, or sulfonic acids. The untreated MDH maintained isoforms clustered around the more basic pI range, as illustrated by the green gel spots 6–7. The relative MDH_{reduction}/MDH_{oxidation} ratios for all MDH isoforms are marked below the corresponding gel spots, with isoform #1 being the most oxidized and isoform #7 being the most reduced. Ratio*: Cy3m/Cy5m ratio represents relative free cysteine thiol levels in control/H₂O₂-treated MDH isoforms.

Table 2. Purified Malate Dehydrogenase Peptide Identification and Thiol-Oxidative Quantification by ICAT

peptide sequence ^a	C.I. %	observed mass	error (ppm)	ICAT ratio (H/L) ^b
<u>⁵⁵G Y L G P E Q L P D C L K</u> ⁶⁷	100	1659.8	-11	0.34
<u>⁶⁸G C D V V V I P A G V P R</u> ⁸⁰	100	1508.8	-13	0.23
<u>¹⁸⁰T I I P L I S Q C T P K</u> ¹⁹¹	100	1540.9	-16	0.49
<u>²⁴⁶E G V V E C S F V K</u> ²⁵⁵	100	1323.5	5	0.45
<u>²⁵⁶S Q E T D C P Y F S T P L L L G K</u> ²⁷²	100	2126.0	-4	0.47

^a ICAT labeled cysteines are underlined. ^b H/L: H₂O₂ oxidized/untreated MDH peptides.

from the tissues, which is important for biological studies. Because in our model system the proteins were oxidized by H₂O₂ after denaturation, many cysteine thiols might not possess their native protein structures in denaturing lysis buffer; therefore protein sensitivities to oxidant H₂O₂ reported in this study may not be identical to *in vivo* conditions. However, because no reduction and alkylation steps were included in the lysis buffer, select cysteine thiols might still contain some structural environments that enabled their interactions with other vicinal amino acids, resulting in varied thiol sensitivity to H₂O₂ oxidation observed in this study (see example in Figure 5). We can not claim unequivocally whether the isolated heart proteins oxidized by 500 μM of H₂O₂ are also sensitive to intracellular oxidants under pathophysiological conditions. However, from two subsequent independent studies of heart proteins isolated from animal models with deficiencies of redox modulators, we found many of the same proteins were differentially oxidized (manuscripts in preparation), confirming the effectiveness of both DIGE and ICAT methods for the discovery of distinct groups of redox-sensitive proteins.

Taken together, both methods have demonstrated the effectiveness for the identification of both previously known and unknown redox-sensitive proteins. Meanwhile, each method was useful at discovering a unique set of novel redox-sensitive proteins, and may pave the way for further mechanistic studies. DIGE could be useful for the identification of redox-induced protein isoform formations and ICAT method is more versatile for identifying redox-sensitive peptides with exact localization of the cysteine oxidized. Overall, these two methods are complementary for redox proteomics studies. The selection of the best method is task-dependent (Table 3). If the goal is to quantify the global redox proteomic changes, DIGE is the

Table 3. Relative Merits of DIGE and ICAT Techniques for Protein Thiol Quantification

	advantages	disadvantages
DIGE	<ul style="list-style-type: none"> • More confident protein identification, better sequence coverage • More effective at identifying protein oxidative isoforms • Easier implementation of statistics for multiplexed quantification • Suitable for large scale screening 	<ul style="list-style-type: none"> • Not amenable for analyzing basic, hydrophobic, and large proteins • Unable to locate the exact sites of cysteine oxidative modifications • Comigration of proteins may interfere with accurate protein identification and thiol quantification • May overlook low-abundant proteins
ICAT	<ul style="list-style-type: none"> • Able to locate the exact sites of redox-sensitive cysteines • Amenable to multidimensional separation and enhances the detection of low-abundant proteins • Suitable for large scale screening 	<ul style="list-style-type: none"> • Less protein sequence coverage than 2DE • Multiple ICAT experiments are required to establish statistical significance for quantification and may be time-consuming • Overlapping ions in MS spectra may interfere with accurate quantification and identification

method of choice. However, if the focus is to identify specific cysteines within key proteins for downstream structural and functional study, ICAT may be more advantageous. If the goal is to identify as many redox-sensitive proteins as possible, both methods may be needed.

Summary

A systematic comparison between DIGE and ICAT methods was carried out for detecting redox-sensitive proteins following the H₂O₂ treatment of heart proteins. Each method complemented the other and can be used together to obtain a comprehensive understanding of the changes in heart redox proteome. ICAT is an attractive tool for the precise localization of redox-sensitive cysteine in low-abundant proteins, whereas DIGE is advantageous for easier implementation of multiplexed experiments to discover redox proteomics patterns. Collectively,

we have identified 63 unique redox-sensitive proteins, including some previously reported and several novel H₂O₂ oxidation targets, that may shed light into better understandings of the redox-mediated regulation of cellular processes.

Acknowledgment. This research is supported in part by a NIH grant NS046593 to H.L. and by a grant from the foundation of UMDNJ to H.L. and J.S.

Supporting Information Available: Supplemental Tables 1–3 and Supplemental Figures 1 and 2. This material is available free of charge via the Internet at <http://pubs.acs.org>.

References

- Ghezzi, P.; Bonetto, V. Redox proteomics: Identification of oxidatively modified proteins. *Proteomics* **2003**, *3* (7), 1145–53.
- Di Simplicio, P.; Franconi, F.; Frosali, S.; Di Giuseppe, D. Thiolation and nitrosation of cysteines in biological fluids and cells. *Amino Acids* **2003**, *25* (3–4), 323–39.
- Zheng, M.; Aslund, F.; Storz, G. Activation of the OxyR transcription factor by reversible disulfide bond formation. *Science* **1998**, *279* (5357), 1718–21.
- Abate, C.; Patel, L.; Rauscher, F. J.; Curran, T. Redox regulation of fos and jun DNA-binding activity in vitro. *Science* **1990**, *249* (4973), 1157–61.
- Toledano, M. B.; Leonard, W. J. Modulation of transcription factor NF-kappa B binding activity by oxidation-reduction in vitro. *Proc. Natl. Acad. Sci. U.S.A.* **1991**, *88* (10), 4328–32.
- Xanthoudakis, S.; Miao, G. G.; Curran, T. The redox and DNA-repair activities of Ref-1 are encoded by nonoverlapping domains. *Proc. Natl. Acad. Sci. U.S.A.* **1994**, *91* (1), 23–7.
- Rainwater, R.; Parks, D.; Anderson, M. E.; Tegtmeyer, P.; Mann, K. Role of cysteine residues in regulation of p53 function. *Mol. Cell. Biol.* **1995**, *15* (7), 3892–903.
- Gu, J.; Milligan, J.; Huang, L. E. Molecular mechanism of hypoxia-inducible factor 1alpha-p300 interaction. A leucine-rich interface regulated by a single cysteine. *J. Biol. Chem.* **2001**, *276* (5), 3550–4.
- Humphries, K. M.; Juliano, C.; Taylor, S. S. Regulation of cAMP-dependent protein kinase activity by glutathionylation. *J. Biol. Chem.* **2002**, *277* (45), 43505–11.
- Kamata, H.; Honda, S.; Maeda, S.; Chang, L.; Hirata, H.; Karin, M. Reactive oxygen species promote TNFalpha-induced death and sustained JNK activation by inhibiting MAP kinase phosphatases. *Cell* **2005**, *120* (5), 649–61.
- Tonks, N. K. Redox redux: revisiting PTPs and the control of cell signaling. *Cell* **2005**, *121* (5), 667–70.
- Park, H. S.; Yu, J. W.; Cho, J. H.; Kim, M. S.; Huh, S. H.; Ryoo, K.; Choi, E. J. Inhibition of apoptosis signal-regulating kinase 1 by nitric oxide through a thiol redox mechanism. *J. Biol. Chem.* **2004**, *279* (9), 7584–90.
- Vivancos, A. P.; Castillo, E. A.; Biteau, B.; Nicot, C.; Ayte, J.; Toledano, M. B.; Hidalgo, E. A cysteine-sulfenic acid in peroxiredoxin regulates H₂O₂-sensing by the antioxidant Pap1 pathway. *Proc. Natl. Acad. Sci. U.S.A.* **2005**, *102* (25), 8875–80.
- Fujiwara, N.; Nakano, M.; Kato, S.; Yoshihara, D.; Ookawara, T.; Eguchi, H.; Taniguchi, N.; Suzuki, K. Oxidative modification to cysteine sulfonic acid of Cys111 in human copper-zinc superoxide dismutase. *J. Biol. Chem.* **2007**, *282* (49), 35933–44.
- Sumbayev, V. V. S-nitrosylation of thioredoxin mediates activation of apoptosis signal-regulating kinase 1. *Arch. Biochem. Biophys.* **2003**, *415* (1), 133–6.
- Graf, P. C.; Jakob, U. Redox-regulated molecular chaperones. *Cell. Mol. Life Sci.* **2002**, *59* (10), 1624–31.
- Ilbert, M.; Horst, J.; Ahrens, S.; Winter, J.; Graf, P. C.; Lilie, H.; Jakob, U. The redox-switch domain of Hsp33 functions as dual stress sensor. *Nat. Struct. Mol. Biol.* **2007**, *14* (6), 556–63.
- Brennan, J. P.; Wait, R.; Begum, S.; Bell, J. R.; Dunn, M. J.; Eaton, P. Detection and mapping of widespread intermolecular protein disulfide formation during cardiac oxidative stress using proteomics with diagonal electrophoresis. *J. Biol. Chem.* **2004**, *279* (40), 41352–41360.
- Charles, R. L.; Schroder, E.; May, G.; Free, P.; Gaffney, P. R.; Wait, R.; Begum, S.; Heads, R. J.; Eaton, P. Protein sulfenation as a redox sensor: proteomics studies using a novel biotinylated dimedone analogue. *Mol. Cell. Proteomics* **2007**, *6* (9), 1473–84.
- Saurin, A. T.; Neubert, H.; Brennan, J. P.; Eaton, P. Widespread sulfenic acid formation in tissues in response to hydrogen peroxide. *Proc. Natl. Acad. Sci. U.S.A.* **2004**, *101* (52), 17982–87.
- Hao, G.; Derakhshan, B.; Shi, L.; Campagne, F.; Gross, S. S. SNOSID, a proteomic method for identification of cysteine S-nitrosylation sites in complex protein mixtures. *Proc. Natl. Acad. Sci. U.S.A.* **2006**, *103* (4), 1012–17.
- Fratelli, M.; Demol, H.; Puype, M.; Casagrande, S.; Eberini, I.; Salmona, M.; Bonetto, V.; Mengozzi, M.; Duffieux, F.; Miclet, E.; Bachi, A.; Vandekerckhove, J.; Gianazza, E.; Ghezzi, P. Identification by redox proteomics of glutathionylated proteins in oxidatively stressed human T lymphocytes. *Proc. Natl. Acad. Sci. U.S.A.* **2002**, *99* (6), 3505–10.
- Michelet, L.; Zaffagnini, M.; Marchand, C.; Collin, V.; Decottignies, P.; Tsan, P.; Lancelin, J. M.; Trost, P.; Miginiac-Maslow, M.; Noctor, G.; Lemaire, S. D. Glutathionylation of chloroplast thioredoxin f is a redox signaling mechanism in plants. *Proc. Natl. Acad. Sci. U.S.A.* **2005**, *102* (45), 16478–83.
- Yano, H.; Wong, J. H.; Lee, Y. M.; Cho, M. J.; Buchanan, B. B. A strategy for the identification of proteins targeted by thioredoxin. *Proc. Natl. Acad. Sci. U.S.A.* **2001**, *98* (8), 4794–99.
- Essex, D. W.; Li, M. R.; Miller, A.; Feinman, R. D. Protein disulfide isomerase and sulfhydryl-dependent pathways in platelet activation. *Biochemistry* **2001**, *40* (20), 6070–75.
- Baty, J. W.; Hampton, M. B.; Winterbourn, C. C. Detection of oxidant sensitive thiol proteins by fluorescence labeling and two-dimensional electrophoresis. *Proteomics* **2002**, *2* (9), 1261–66.
- Kim, J. R.; Yoon, H. W.; Kwon, K. S.; Lee, S. R.; Rhee, S. G. Identification of proteins containing cysteine residues that are sensitive to oxidation by hydrogen peroxide at neutral pH. *Anal. Biochem.* **2000**, *283* (2), 214–21.
- Woo, H. A.; Kang, S. W.; Kim, H. K.; Yang, K. S.; Chae, H. Z.; Rhee, S. G. Reversible oxidation of the active site cysteine of peroxiredoxins to cysteine sulfenic acid. Immunoblot detection with antibodies specific for the hyperoxidized cysteine-containing sequence. *J. Biol. Chem.* **2003**, *278* (48), 47361–4.
- Leichert, L. I.; Jakob, U. Protein thiol modifications visualized in vivo. *PLoS Biol.* **2004**, *2* (11), e333.
- Hurd, T. R.; Prime, T. A.; Harbour, M. E.; Lilley, K. S.; Murphy, M. P. Detection of ros-sensitive thiol proteins by redox-difference gel electrophoresis (redox-dige): Implications for mitochondrial redox signalling. *J. Biol. Chem.* **2007**, 22040–51.
- Gygi, S. P.; Rist, B.; Gerber, S. A.; Turecek, F.; Gelb, M. H.; Aebersold, R. Quantitative analysis of complex protein mixtures using isotope-coded affinity tags. *Nat. Biotechnol.* **1999**, *17* (10), 994–9.
- Sethuraman, M.; McComb, M. E.; Huang, H.; Huang, S. Q.; Heibeck, T.; Costello, C. E.; Cohen, R. A. Isotope-coded affinity tag (ICAT) approach to redox proteomics: Identification and quantitation of oxidant-sensitive cysteine thiols in complex protein mixtures. *J. Proteome Res.* **2004**, *3* (6), 1228–33.
- Christoffers, K. H.; Li, H.; Howells, R. D. Purification and mass spectrometric analysis of the delta opioid receptor. *Mol. Brain Res.* **2005**, *136* (1–2), 54–64.
- Liu, T.; Donahue, K. C.; Hu, J.; Kurnellas, M. P.; Grant, J. E.; Li, H.; Elkabes, S. Identification of differentially expressed proteins in experimental autoimmune encephalomyelitis (EAE) by proteomic analysis of the spinal cord. *J. Proteome Res.* **2007**, *6* (7), 2565–75.
- Peng, J.; Elias, J. E.; Thoreen, C. C.; Licklider, L. J.; Gygi, S. P. Evaluation of multidimensional chromatography coupled with tandem mass spectrometry (LC/LC-MS/MS) for large-scale protein analysis: the yeast proteome. *J. Proteome Res.* **2003**, *2* (1), 43–50.
- Lee, C. K.; Park, H. J.; So, H. H.; Kim, H. J.; Lee, K. S.; Choi, W. S.; Lee, H. M.; Won, K. J.; Yoon, T. J.; Park, T. K.; Kim, B. Proteomic profiling and identification of cofilin responding to oxidative stress in vascular smooth muscle. *Proteomics* **2006**, *6* (24), 6455–75.
- Obata, T. Nitric oxide and depolarization induce hydroxyl radical generation. *Jpn. J. Pharmacol.* **2002**, *88* (1), 1–5.
- Yang, K.-S.; Kang, S. W.; Woo, H. A.; Hwang, S. C.; Chae, H. Z.; Kim, K.; Rhee, S. G. Inactivation of Human Peroxiredoxin I during Catalysis as the Result of the Oxidation of the Catalytic Site Cysteine to Cysteine-sulfenic Acid. *J. Biol. Chem.* **2002**, *277* (41), 38029–36.
- Canet-Aviles, R. M.; Wilson, M. A.; Miller, D. W.; Ahmad, R.; McLendon, C.; Bandyopadhyay, S.; Baptista, M. J.; Ringe, D.; Petsko, G. A.; Cookson, M. R. The Parkinson's disease protein DJ-1 is neuroprotective due to cysteine-sulfenic acid-driven mitochondrial localization. *Proc. Natl. Acad. Sci. U.S.A.* **2004**, *101* (24), 9103–8.

- (40) Janssen-Heininger, Y. M.; Poynter, M. E.; Baeuerle, P. A. Recent advances towards understanding redox mechanisms in the activation of nuclear factor kappaB. *Free Radic. Biol. Med.* **2000**, *28* (9), 1317–27.
- (41) Sethuraman, M.; McComb, M. E.; Heibeck, T.; Costello, C. E.; Cohen, R. A. Isotope-coded affinity tag approach to identify and quantify oxidant-sensitive protein thiols. *Mol. Cell. Proteomics* **2004**, *3* (3), 273–8.
- (42) Li, Y.; Camacho, P. Ca²⁺-dependent redox modulation of SERCA 2b by ERp57. *J. Cell Biol.* **2004**, *164* (1), 35–46.
- (43) Declercq, J. P.; Evrard, C.; Clippe, A.; Stricht, D. V.; Bernard, A.; Knoops, B. Crystal structure of human peroxiredoxin 5, a novel type of mammalian peroxiredoxin at 1.5 Å resolution. *J. Mol. Biol.* **2001**, *311* (4), 751–9.
- (44) Nakajima, H.; Amano, W.; Fujita, A.; Fukuhara, A.; Azuma, Y.-T.; Hata, F.; Inui, T.; Takeuchi, T. The Active Site Cysteine of the Proapoptotic Protein Glyceraldehyde-3-phosphate Dehydrogenase Is Essential in Oxidative Stress-induced Aggregation and Cell Death. *J. Biol. Chem.* **2007**, *282* (36), 26562–74.
- (45) Wolosker, H.; Panizzutti, R.; Engelender, S. Inhibition of creatine kinase by S-nitrosoglutathione. *FEBS Lett.* **1996**, *392* (3), 274–6.
- (46) Sharma, A. B.; Sun, J.; Howard, L. L.; Williams, A. G., Jr.; Mallet, R. T. Oxidative stress reversibly inactivates myocardial enzymes during cardiac arrest. *Am. J. Physiol. Heart Circ. Physiol.* **2007**, *292* (1), H198–206.
- (47) Inarrea, P.; Moini, H.; Rettori, D.; Han, D.; Martinez, J.; Garcia, I.; Fernandez-Vizarra, E.; Iturralde, M.; Cadenas, E. Redox activation of mitochondrial intermembrane space Cu,Zn-superoxide dismutase. *Biochem. J.* **2005**, *387* (Pt 1), 203–9.
- (48) Han, D.; Canali, R.; Garcia, J.; Aguilera, R.; Gallaher, T. K.; Cadenas, E. Sites and mechanisms of aconitase inactivation by peroxynitrite: modulation by citrate and glutathione. *Biochemistry* **2005**, *44* (36), 11986–96.
- (49) Eaton, P.; Byers, H. L.; Leeds, N.; Ward, M. A.; Shattock, M. J. Detection, quantitation, purification, and identification of cardiac proteins S-thiolated during ischemia and reperfusion. *J. Biol. Chem.* **2002**, *277* (12), 9806–11.
- (50) Cabiscol, E.; Piulats, E.; Echave, P.; Herrero, E.; Ros, J. Oxidative stress promotes specific protein damage in *Saccharomyces cerevisiae*. *J. Biol. Chem.* **2000**, *275* (35), 27393–8.
- (51) Chen, Y. R.; Chen, C. L.; Pfeiffer, D. R.; Zweier, J. L. Mitochondrial complex II in the post-ischemic heart: oxidative injury and the role of protein S-glutathionylation. *J. Biol. Chem.* **2007**, *282* (45), 32640–54.
- (52) Meany, D. L.; Xie, H.; Thompson, L. V.; Arriaga, E. A.; Griffin, T. J. Identification of carbonylated proteins from enriched rat skeletal muscle mitochondria using affinity chromatography-stable isotope labeling and tandem mass spectrometry. *Proteomics* **2007**, *7* (7), 1150–63.
- (53) Newman, S. F.; Sultana, R.; Perluigi, M.; Coccia, R.; Cai, J.; Pierce, W. M.; Klein, J. B.; Turner, D. M.; Butterfield, D. A. An increase in S-glutathionylated proteins in the Alzheimer's disease inferior parietal lobule, a proteomics approach. *J. Neurosci. Res.* **2007**, *85* (7), 1506–14.
- (54) Schilling, B.; Yoo, C. B.; Collins, C. J.; Gibson, B. W. Determining cysteine oxidation status using differential alkylation. *Int. J. Mass Spectrom.* **2004**, *236* (1–3), 117–27.
- (55) Mohr, S.; Stamler, J. S.; Brune, B. Posttranslational modification of glyceraldehyde-3-phosphate dehydrogenase by S-nitrosylation and subsequent NADH attachment. *J. Biol. Chem.* **1996**, *271* (8), 4209–14.
- (56) Ishii, T.; Tatsuda, E.; Kumazawa, S.; Nakayama, T.; Uchida, K. Molecular basis of enzyme inactivation by an endogenous electrophile 4-hydroxy-2-nonenal: identification of modification sites in glyceraldehyde-3-phosphate dehydrogenase. *Biochemistry* **2003**, *42* (12), 3474–80.
- (57) Kim, B. J.; Hood, B. L.; Aragon, R. A.; Hardwick, J. P.; Conrads, T. P.; Veenstra, T. D.; Song, B. J. Increased oxidation and degradation of cytosolic proteins in alcohol-exposed mouse liver and hepatoma cells. *Proteomics* **2006**, *6* (4), 1250–60.
- (58) Nagumo, Y.; Kakeya, H.; Shoji, M.; Hayashi, Y.; Dohmae, N.; Osada, H. Epolactaene binds human Hsp60 Cys442 resulting in the inhibition of chaperone activity. *Biochem. J.* **2005**, *387* (Pt 3), 835–40.

PR800233R