
Know your meds



- Albuterol Pumps (aka Metered dose inhalers- MDIs)
- Nebulizer medication
 - Albuterol
 - Inhaled steroids
 - Other
- o Special albuterol forms
- Inhaled Steroids
- Inhaled Steroids- Dry Powder Inhalers (DPIs)
- o Combo medications- LABA/ICS MDIs
- o Combo medications- LABA/ ICS DPIs
- o Asthma Shots
- Other Medications- Common in different ages and diseases
- o Look- A- Likes

Albuterol Pumps













Nebulized Albuterol







Nebulized Inhaled Steroids







Nebulized Inhaled Steroids





Nebulized Inhaled Steroids









Nebulized others



30 mL

NOT FOR INJECTION

0 Solution, USP

For Inhalation (Mucolytic Agent) or Oral Administration (Acetaminophen Antidote).

Distributed by Hospira, Inc., Lake Forest, IL 60045 USA

##-###-AA

DMMMYYYY

NDC 0409-3307-11 Rx only

Each mL contains acetylcysteine 100 mg; edetate disodium, dihyd. 0.25 Contains NaOH and may contain HČI for pH adjustment. Usual Dosage: See accompanying directions for dosage, indications and use. Store at 20 to 25°C (68 to 77°F). [See USP Controlled Room Temperature.] Store in refrigerator at 2° to 8°C (36° to 46°F) after opening. 2° to 8°C (36° to 46°F) after opening. Discard opened vial after 96 hours. A change in color may occur after opening; this does properly in change the efficacy of the drug.

RL - 7544 Hośpira not change the efficacy of the drug.





NDC 63323-690-30

ACETYLCYSTEINE

SOLUTION, USP

20%

(200 mg/mL)

For Inhalation (Mucolytic Agent) or Oral Administration (Acetaminophen Antidote)

NOT FOR INJECTION, PRESERVATIVE FREE

Each ml. contains 200 mg (20%) Acetylcysteine, 0.05% disodium edetate, sodium hydroxide and, if necessary, hydrochloric acid is added (used for pH adjustment) and water for injection. pH (range 6.0 to 7.5) Usual Dosage: See Insert.

WARNING: DISCARD OPENED CONTAINERS AFTER 96 HOURS

Store at 20°: 25°C (68°-77°F); excursions permitted to 15°-30°C (59°-86°F) [see USP Controlled Room Temperature]. Store in refrigerator 2* - 8*C (36* - 46*F) after opening. A change in color may occur after opening. This does not change the efficacy of the drug. Acetylcysteine may be diluted to a lesser concentration with an appropriate solution.

Manufactured for:



FRESENIUS Lake Zurich, IL 60047 Made in India

www.fresenius-kabi.com/us

Sample Label. Please see package insert for complete

prescribing


information.

ode No.: AP/DRUGS/103/97 315000008-02


3 x 30 mL Vials

Rx only

Neb others (cont)









Special meds









Inhaled Steroids













Asmanex® HFA
(mometasone furoate)
Inhalation Aerosol

200 mcg per actuation
For oral inhalation only
Asmanex HFA canister to be used with Asmanex HFA actuator only.

SHAKE WELL BEFORE USING.

Rx only

120 Metered Actuations
Net Wt. 139



Inhaled steroids- Dry powder











00186-0916-12 PULMICORT FLX 180 MCG INH 120 AEROSOL , WHITE

00186-0917-06 PULMICORT FLX 90 MCG INH 60

Combo Meds LABA ICS






















NDC 0085-4610-01


LABA ICS DPI












100 mcg/25 mcg









Asthma Shots









Asthma shots(cont)









er medications





20 mcg/100 mcg per actuation*

FOR ORAL INHALATION ONLY

4 Grams 120 Metered Doses

Boehringer Ingelheim











Aqua (2.5 mcg/puff)

Blue (1.25 mcg/puff)


IIII

ms Combivent- Respirat

FOR ORAL DOMALATION ON

Ohner.

Look A-likes







NDC 78206-125-01

Dulera°

I Dulera

Dulera*

inconstance funcable

Inhalation Aerosof

50 mcg/5 mcg per actuation for oral inhabition only

Use only with Dolera® canister. Do not remove

SHARE WELL BEFORE USING.

ST THE RESIDENCE

7006245800

(mometasone furoate and

Aerosol

EFORE USING.

Net Wt. 13g









Look A likes















Look A likes








Look A Likes





Look A Likes





NOT TRADITIONAL ASTHMA MEDICATION







Primatene®



